Fayetteville Gastroenterology Associates 2041 Valleygate Dr. Fayetteville, NC 28304 (910) 323-5203

Please report to the second floor at	800
Your procedure will follow a brief pre-op interview.	

Split Dose Colonoscopy Prep with Miralax

	You will need to purchase Miralax gram bottle over the counter.
	This is the most important part of your colonoscopy. A clean colon provides optimal evaluation. If your colon is not clean, polyps and cancers may be missed.
	The Day Before Your Procedure
	You will be on a clear liquid diet the entire day. See the back of this page for specific instructions.
	Early in the day, mix the entire container of Miralax with 64 ounces of Gatorade o Do not use red or purple Gatorade o Gatorade should be room temperature to mix, but once the Miralax powder is added and dissolves completely, the solution can be refrigerated
	At take 4 Dulcolax pills with 8oz. of water.
	At begin drinking the Miralax solution. Drink an 8oz glass every 20-25 minutes until 2/3 of the container is gone. o Do not sip! o Initially you may feel bloated, but you will feel better once you begin to have bowel movements. o Put the remaining solution back in the refrigerator.
	The Day of Your Procedure
	At begin drinking the remainder of the solution, an 8 oz glass every 20-25 minutes, until the container is empty.
	You may continue clear liquids until 2 hours prior to your appointment time. You should have nothing by mouth after This includes water, gum, hard candy, cough drops, and medications.
If yo	ou have anything by mouth within 2 hours of your appointment, your procedure may be cancelled.
	If you have any questions during normal business hours,
	please call at
	If you need assistance after normal business hours

If you need assistance after normal business hours (after 4:30pm Monday-Thursday, 1pm on Friday, or on the weekend) Please call (910) 323-5203 and press option 8 for the answering service.